

Atraumatic Restorative Treatment (ART) as Public Policy: a Scoping Review

O Tratamento Restaurador Atraumático (ART) como Política Pública: uma Revisão de Escopo

Fábio Carneiro Martins^{*ab}; Laura Regina Antunes Pontes^a; Gabriela da Silva Martins Gonçalves^a; Mariana Gabriel^a; Daniela Prócida Raggio^a; Mariana Minatel Braga^a; Antonio Carlos Frias^a; Fernanda Campos de Almeida Carrer^a

^aUniversidade de São Paulo, Faculty of Dentistry, Department of Community Dentistry, School of Dentistry. SP, Brazil;

^bUniversidade de Brasília, School of Health Sciences. DF, Brazil.

*E-mail: fabio.carneiro.martins@usp.br.

Abstract

This scoping review aimed to identify, describe, and analyze the use of Atraumatic Restorative Treatment (ART) as public policy. Inclusion criteria: studies on public oral health policies; dental caries; ART. Exclusion criteria: clinical studies; specific interventions; studies that report public policies that do not use ART as a strategy for caries treatment. The search was conducted in PubMed, BVS, Epistemonikos, Health Systems Evidence, and Rxforchange. The selection process and data extraction were performed by two authors independently according to the stages and categories of a conceptual framework. Of the 2253 eligible articles, 139 were duplicates, and 1680 were not included after reading the title and abstract. After assessing the full text, 414 articles were excluded. Twenty articles were included. Nineteen were conducted in nine countries (South Africa, Bolivia, Cambodia, Egypt, Mexico, Tanzania, East Timor, Tunisia, and Zimbabwe), and one in the Americas. The studies reported high prevalence of dental caries and lack of access to restorative procedures in health services as common problems. Policy developments followed a similar process, with the presence of governments, conceptual use of scientific evidence, and induction by the World Health Organization. Regarding implementation barriers, the included studies mentioned the lack of supplies and lack of induction by managers, while in relation to the facilitators, permanent education and professional practice were mentioned. Cohort studies have shown a survival rate greater than 80% after one year of follow up of the restorations performed. The findings of this review indicate that the use of ART in public policies is promising, however, its use is still in an early stage.

Keywords: Dental Atraumatic Restorative Treatment. Dental Caries. Public Policy. Community Dentistry. Oral Health.

Resumo

Esta revisão de escopo teve como objetivo identificar, descrever e analisar o uso do Tratamento Restaurador Atraumático (ART) como política pública. Critérios de inclusão: estudos sobre políticas públicas de saúde bucal; cáries dentárias; ART. Critérios de exclusão: estudos clínicos; intervenções específicas; estudos que relatam políticas públicas que não utilizam o ART como estratégia para o tratamento de cárie. A pesquisa foi realizada no PubMed, BVS, Epistemonikos, Health Systems Evidence e Rxforchange. O processo de seleção e extração de dados foram realizados por dois autores, de forma independente, de acordo com as etapas e categorias de um quadro conceitual. Dos 2.253 artigos elegíveis, 139 eram duplicados e 1.680 não foram incluídos após a leitura do título e do resumo. Após avaliação do texto completo, foram excluídos 414 artigos. Vinte artigos foram incluídos. Dezenove foram realizados em nove países (África do Sul, Bolívia, Camboja, Egito, México, Tanzânia, Timor Leste, Tunísia e Zimbábue) e um nas Américas. Os estudos relataram alta prevalência de cárie dentária e falta de acesso a procedimentos restauradores nos serviços de saúde como problemas comuns. Os desenvolvimentos de políticas seguiram um processo semelhante, com a presença de governos, uso conceitual de evidências científicas e indução da Organização Mundial da Saúde. Quanto às barreiras de implementação, foram mencionadas a falta de insumos e a falta de indução por parte dos gestores. Quanto aos facilitadores, foram citados a educação permanente e a prática profissional. Estudos de coorte mostraram sobrevida superior a 80% após um ano de acompanhamento. Os achados desta revisão indicam que a utilização da ART nas políticas públicas é promissora, porém, sua utilização ainda é incipiente.

Palavras-chave: Tratamento Dentário Restaurador sem Trauma. Cárie Dentária. Política Pública. Odontologia Comunitária. Saúde Bucal.

1 Introduction

Illness is a complex phenomenon that needs to be understood in terms of its biopsychosocial dimensions. Actions, programs, or policies that wish to interfere with the determinants of the health illness process must contemplate all these aspects. It is worth mentioning that since the Declaration of Alma-Ata in 1978, strategies to combat inequalities and achieve universal health coverage for the world population have been discussed^{1,2}.

Dental caries is a highly prevalent disease in all age groups. Therefore, targeted policies and programs that aim to decrease its prevalence in the population and the health inequities observed in its distribution are required. It is the most common irreversible chronic disease in childhood and cannot be healed by short-term pharmacological interventions³. There is also a lack of equitable public policies that guarantee access to oral health promotion, prevention, cure, and rehabilitation care to those who need it most since oral health is still considered less important in many health care systems around the world⁴.

The Collaborating Center of the World Health Organization (WHO) in Nijmegen published in 2002 a document titled “The Basic Package of Oral Care” (BPOC)⁵. This reference proposes strategies to deal with most oral health needs of a population in primary health care. Atraumatic Restorative Treatment (ART) is a potential treatment.

ART was conceived by Joe E. Frencken and collaborators in the mid-1980s in Tanzania to be conducted in social spaces where there is no electricity, which made conventional dental treatments impossible. ART showed high quality and reliability for treating dental caries injuries, indicating that the technique should not be restricted to territories with these characteristics^{6,7}. The ART approach has been widely disseminated worldwide by the WHO, which has recognized it as an appropriate and potentially impactful treatment since 1994.

Currently, ART is understood as a minimally invasive approach, which includes preventive, therapeutic, and restorative measures for dental caries⁸, and is perfectly inserted in the modern philosophy of health care, which is characterized by maximum effort in health-promoting, preventive approaches and in carrying out, when necessary, minimally invasive procedures. In addition, its quality is supported by current and robust scientific evidence^{9,10}, as over the past decades, high quality research on ART has been developed, and several research groups have advocated its implementation in routine health services.

However, despite the intense scientific production on ART, little is known about its use for decision making by managers and the consequent inclusion in public oral health policies. Therefore, the objective of this scoping review was to identify studies that involve the use of ART as a public policy or government program to better understand the use of this technique in routine health care in municipalities, states, or countries around the world.

2 Material and Methods

2.1 Conceptual framework on the theme studied

To better conduct this scoping review, we used two

frameworks previously published: the first was the “3-i Framework: Interests, Ideas and Institutions” developed by François-Pierre Gauvin, from the National Center of Collaboration for Healthy Public Policies in Montréal, Québec¹¹, and the second was a framework designed for analyzing sustainable implementations in oral health, developed by D. Dwayne Simpson, from the Institute of Behavioral Research, Texas Christian University¹².

The first framework brings together three of the most common factors that political science literature uses to explain public policy development processes. Commonly referred to as the “3-i framework,” it declares that development and policy choices are influenced by interests, which are defined by agendas of social groups, such as civil servants, managers, researchers, and politicians; ideas, which include scientific evidence, professional experience, dominant values and culture; and institutions, composed of formal and informal rules, norms, precedents and organizational factors that structure political behavior, that is, government structures, policy networks, and political legacies. The model was recently updated with external factors (such as political changes and the media) as a fourth aspect that also influences the decision making process¹³.

The second framework, a framework for implementing sustainable oral health promotion interventions, focuses on four stages of implementing interventions, which show how the concepts present in each of them contribute as part of an integrated chain of events of key factors that sequentially influence the sustainability of an innovation/intervention in oral health¹². In addition, it highlights the organization, readiness, and infrastructure of health systems as essential aspects for an implementation to be possible and that the availability of resources (financial, human, logistical) added to the management of organizational tensions will impact their sustainability.

From reading the two frameworks described, we synthesized the necessary elements in Chart 1, which was used as the guiding thread of this work

Table 1 - Guiding synthesis for description and analysis of the studied interventions

Stage	Categories	Relevant Aspects	Framework
Development	Context, decision making process, formulation	Problem definition Institutions (Government structures; policy networks; political legacies) Ideas (Scientific knowledge/evidence; dominant values/culture in society) Interests (agenda of social groups, civil servants, researchers, and politicians)	3-I + E
Implementation	Permanent education	Regarding pilot projects: importance of leadership, flexibility, usefulness, adaptability, quality Regarding courses: compatibility with the needs of professionals, accessibility, certification offer	A framework for implementing sustainable oral health promotion interventions
	Data on the implementation process and its impact (quantitative and qualitative data)	Quantity and longevity of restorative procedures performed Facilitators (Effectiveness, feasibility, versatility, availability of inputs, targets, monitoring, permanent education) Barriers (turnover in management and health teams, resistance, low fidelity)	A framework for implementing sustainable oral health promotion interventions

Source: adapted from the Frameworks¹¹

2.2 Eligibility criteria

A broad research question was formulated: What is the current global scenario described in the scientific literature regarding the use of ART as a public policy?

It is worth mentioning that there is no single or better definition of what public policy is, but there are different meanings that guide the discussion to the locus where the clashes of interests, preferences, and ideas develop, that is, in governments. Therefore, we will consider public policy as the process by which governments translate their purposes into actions in search of results or changes in a known reality¹⁴.

Regarding selection, there were no restrictions on the date of publication or language.

Inclusion criteria: Studies on public oral health policies, dental caries, or ART. Exclusion criteria: Clinical studies, specific interventions unrelated to the decision making of

managers or policymakers, and studies that report public oral health policies that do not use ART as a coping strategy for dental caries.

2.3 Information sources and search strategies

A literature search was conducted on PubMed, Biblioteca Virtual em Saúde- BVS, Epistemonikos, Health Systems Evidence, Rx for change, and the Cochrane Library databases. To complete the search, we sent an invitation via email so that key informants (researchers who participated in the idealization and implementation of ART in research programs or projects) could report experiences via a form created with the *Google Forms* tool. The last search was conducted on March 5, 2021.

The details of the search strategies used are described in Table 2.

Table 2 - Search strategy

Database	Search	Nº Articles Found
PubMed	(((((("oral care"[Title/Abstract] OR "salud oral"[Title/Abstract]) OR "oral health care"[Title/Abstract]) OR "oral health"[MeSH Terms]) OR "oral health"[Title/Abstract]) OR "public policy"[MeSH Terms]) OR "public policy"[All Fields]) OR "community dentistry"[All Fields]) OR "government"[Title/Abstract] OR "public health"[Title/Abstract] OR "health system"[Title/Abstract]) AND (((("atraumatic restorative treatment"[Title/Abstract] OR "dental treatment"[Title/Abstract]) OR "ART approach"[Title/Abstract]) OR "Atraumatic Restorative Treatment (ART)"[Title/Abstract]) OR "ART restorations"[Title/Abstract]) OR "dental atraumatic restorative treatment"[MeSH Terms])	2048
BVS	tw:((tw:(public policy)) OR (tw:(oral health)) AND (tw:(atraumatic restorative treatment)))	187
Epistemonikos	(title:(title:(oral health) OR abstract:(oral health)) OR (title:(oral care) OR abstract:(oral care)) AND (title:(atraumatic restorative treatment) OR abstract:(atraumatic restorative treatment))) OR abstract:(title:(oral health) OR abstract:(oral health)) OR (title:(oral care) OR abstract:(oral care)) AND (title:(atraumatic restorative treatment) OR abstract:(atraumatic restorative treatment)))	14
Health Systems Evidence	Public Policy AND dental caries	1
Rx for change	Public Policy AND Atraumatic Restorative Treatment	2
Other sources (Key informants)	Not applicable	1
Total		2253

Source: research data.

After implementing the search strategies, 2253 studies were screened.

2.4 Studies selection

The titles and abstracts of the articles selected in the electronic search were exported from the databases and organized in Microsoft Excel 365. Two reviewers (FCM; GSMG) performed the calibration with 10% of the sample in the initial search (n = 206); Cohen Kappa's coefficient was 0.81. The inclusion and exclusion phases occurred independently, and in cases of disagreement, we reached a consensus. A third reviewer did not need to participate.

2.5 Data collection and Risk of bias

The extraction of qualitative and quantitative data

of interest was conducted by two authors independently (FCM; LRAP), according to the stages and categories of the conceptual framework (Chart 1), and organized in spreadsheets using Microsoft Excel 365. The data were also described and analyzed from the same perspective, stages, and categories.

We combined the results of qualitative data related to the "development" stages. The experiences were subdivided into the following categories: a) Experiences that stopped in the introduction stage were not implemented or were not consolidated as a policy; b) Unstable experiences; c) Sustainable experiences.


No risk of bias analysis was performed. This review aimed to synthesize, contextualize, describe, and subjectively analyze the experiences of countries rather than evaluate the design of the studies.

3 Results and Discussion

3.1 Selection and general characteristics of the studies

Figure 1 illustrates the study selection process. Of the 2253 articles selected through the database and key informants. After removing 139 duplicates, 1680 were excluded based on the inclusion criteria and 414 by the exclusion criteria.

Figure 1 - Flow Diagram



Source: adapted from PRISMA flow gram¹⁵

Twenty studies were selected, of which nineteen were conducted in nine countries, South Africa (n = 3)¹⁶⁻¹⁸, Bolivia (n = 1)¹⁹, Cambodia (n = 2)^{20,21}, Egypt (n = 2)^{22,23}, Mexico (n = 2)^{24,25}, Tanzania (n = 4)²⁶⁻²⁹, East Timor (n = 1)³⁰, Tunisia (n = 1)³¹, and Zimbabwe (n = 3)³²⁻³⁴ and the other one contained data on the integration of ART in the health systems of countries in the Americas³⁵, based on the opinions of key informants.

As for the year of publication, studies from 1996 to 2016 were included. Nineteen studies were published in English, and one in Spanish. Case studies and experience reports (n = 3), cross-sectional studies (n = 5), cohort studies (n = 8), case-control studies (n = 2), one narrative literature review (n = 1), and one thesis (n = 1) were included.

Nine experiences that evaluated ART implementation steps as public policy were identified and reported by these 20 articles. It is not known whether the technique has been used (in specific actions and research projects, without the involvement of decision makers for inclusion in the political agenda, or if the strategy is used as a policy but without evaluation and publication of the results in the scientific literature), in addition to the experiences presented, managers affirm that ART is included in national oral health policies in countries such as Brazil, Chile, Ecuador, Peru, Uruguay, and in the agenda of dentistry schools in countries such as Argentina, Brazil, Ecuador, Peru, and Venezuela³⁵.

3.2 Stage 1 - Development

Regarding the definition of the problem, the studies described the following: high prevalence of dental caries in 100% of the countries; lack of access to restorative services in eight of the nine countries; inequities related to access (lack of human resources in remote areas, rural or peripheral to urban centers), in 100% of countries, and performing mutilating procedures (preventable tooth extractions) in seven of the nine countries. Therefore, studies share common problems, such as the high prevalence of dental caries and the lack of access to restorative procedures in health services, aggravated by inequities in access, which reinforces social differences. Another problem recognized by the scientific literature is the difficulty in moving users to perform treatments for caries lesions with shallow and medium cavities when there is still no report of pain. The scientific literature mentions dental consultations scheduled when patients study or work, fear of losing their jobs, misery, and lack of information as the primary causes of absenteeism³⁶⁻³⁸. The late treatment of oral health problems results in unnecessary and preventable treatments, as long as there are strategies for situational diagnosis and early interventions, in addition to measures to promote health and prevent these problems.

ART, designed to be used in places with no access to conventional procedures, is an essential and safe approach to the problems identified by the studies included in this review, as it can be performed in alternative social spaces, such as schools, in collective actions. It seems to have a lower cost than the conventional dental practice⁷. In addition, it is within the scope of minimal intervention dentistry, which focuses on maintaining oral health throughout the life cycle through preventive and minimally invasive care^{8,39}.

Regarding the institutions, as described in Figure 2, the studies report that in most countries (77.7%), government structures were present, with the involvement of their ministries of health or local coordination in the development stage. However, few studies have reported political networks (11.1%) and political legacies (44.4%). There were government structures reported in most countries, with their ministries of health or local coordination in the formulation stage. However, while the studies cite the presence of institutions, few describe their role in decision making and planning for implementation. In addition, few studies have reported on networks and political legacies. Therefore, little is known about the intrinsic characteristics of public oral health programs or policies in which ART is inserted as a routine health service.

Figure 2 - Development: local context and decision-making process

Country	Institutions			Ideas	Interests				External Factors	Sources of consultation
	Government structures	Policy networks	Political legacies	Scientific knowledge/evidence	Social groups	Public service workers	Re-searchers	Politicians	WHO induction	
Experiences that stopped (or are still) in the development stage										
Egypt	X			X		X	X	X	X	Introduction of the ART approach in Egypt: intentions, clinical effects, and perceived barriers.
East Timor				X			X		X	Opportunities in oral health policy for East Timor
n (%)	1 (50)	0	0	2 (100)		1 (50)	2 (100)	1 (50)	2 (100)	
Unstable experiences										
South Africa				X			X		X	Implementation of the ART approach in South Africa: an activity report.
Tanzania	X			X			X		X	Atraumatic restorative treatment (ART): the Tanzanian experience.
Tunisia	X		X	X			X	X	X	Atraumatic Restorative Treatment and glass ionomer sealants in Tunisian children: survival after 3 years.
Zimbabwe	X			X			X	X	X	Atraumatic Restorative Treatment and glass ionomer sealants in a school oral health program in Zimbabwe: evaluation after 1 year.
n (%)	3 (75)	0	1 (25)	4 (100)	0	0	4 (100)	2 (50)	4 (100)	
Sustainable experiences										
Bolivia	X		X	X	X		X	X	X	El tratamiento restaurador atraumático (TRA) como parte de un paquete básico de servicios de salud oral en el área rural de Bolivia.
Cambodia	X		X	X		X	X	X	X	Dental nurse training in Cambodia - a new approach.
Mexico	X	X	X	X			X	X	X	Implementation and preliminary evaluation of an ART strategy in Mexico: a country example.
n (%)	3 (100)	1 (33.3)	3 (100)	3 (100)	1 (33.3)	1 (33.3)	3 (100)	3 (100)	3 (100)	
Total N (%)	7 (77.7)	1 (11.1)	4 (44.4)	9 (100)	1 (11.1)	2(22.2)	9 (100)	6 (66.6)	9 (100)	

Source: adapted from the Frameworks¹¹⁻¹³

Regarding ideas, all studies report the influence of scientific evidence or the knowledge of researchers, but the type of evidence used is unclear. Conceptual use of scientific evidence seems to have occurred generally (to provide a general understanding of a given topic), rather than instrumentally use (with specific intentions of using

scientific evidence to solve well-defined problems)⁴⁰. It is worth highlighting that since the 1990s the WHO has widely disseminated the results of research through the production of materials that translated the scientific evidence produced^{5,41}, as an important external inducer of the use of ART, having been reported in all experiments.

The instrumental use of scientific evidence can contribute to developing more solid and sustainable policies in the long term. Still, there are several reasons for underuse, such as 1. The evidence competes with other factors in the policy-making process; 2. the evidence is not valued; 3. the evidence is not relevant; 4. the evidence is not easy to use (the results were not effectively communicated or were not available when decision makers needed it, and 5. decision makers do not have mechanisms to facilitate the use of evidence)^{42,43}.

Finally, regarding interests, in all countries, there were reports of interest by researchers and politicians. However, there is also a lack of information about the role of these actors in decision making processes. Bolivia was the only country where the presence of social groups, such as schoolteachers and the children's families, were reported in relation to the implementation of ART as an intersectoral policy, with health education, supervised toothbrushing, and restorative treatments in the school environment¹⁹. This seems to explain the weakness observed in most ART implementations as a policy. That is, as research related to the instrumental use of evidence for well-informed decision making argues for the participation of representatives of all actors involved in this process, as happens, for example, in deliberative dialog and the synthesizing of scientific evidence, which offer feasible political options for facing well-defined problems⁴⁴.

3.3 Stage 2 - Implementation

3.3.1 Permanent education

Regarding continuing education, we identified the absence of pilot projects in all interventions. According to Chart 1, the aspects that guarantee the effectiveness of preparatory courses and constant updating, as well as the professionals' adherence to the activities, are accessibility (reported in South Africa, Cambodia, Mexico), which involves the release of professionals to attend training, and compatibility with professionals' needs (reported in Cambodia, Mexico), focusing on the work process according to their settings, and certification offer (reported in South Africa), which guarantees professional career progression.

Another study described the same component in a particular way to assess the perception of policy makers in American countries³⁵, using a questionnaire. It reported that all respondents (Chile, Ecuador, El Salvador, Honduras, Mexico, Nicaragua, Panama, Paraguay, Peru, Uruguay, and the United States) claimed to have had courses in their countries and that in 88.2% of those countries, at least two opportunities. The authors reported that ART is included in the agenda of dental schools in Argentina, Bolivia, Brazil, Ecuador, Mexico, Peru, and Venezuela.

The training was described with a different focus (dissemination to large-scale public service professionals was not reported) in two other countries. In Tanzania²⁶, training was provided by universities or external institutions. In Zimbabwe³², only professionals who participated in research projects were involved and performed follow up of the procedures.

3.3.2 Quantitative data inserir também a discussão

Five cohort studies monitored ART restoration survival as described in Table 1.

Table 1 - Quantitative data - restoration survival rates

Studies that follow the ART restorations in single surfaces: Primary teeth							
Study title and year of publication	Country	Total procedures performedn	Sample loss %	Evaluation criteria	Survival rate after one year of follow up %	Survival rate after two years of follow up %	Survival rate after two to three years of follow up %
Atraumatic restorative treatment and glass-ionomer sealants in a school oral health programme in Zimbabwe: evaluation after 1 year (1996)	Zimbabwe	316	32.6	ART codes and criteria for assessing restorations and sealants/criteria for diagnosing carious lesions in ART studies	93.4		
Three-year survival of one-surface ART restorations and glass-ionomer sealants in a school oral health programme in Zimbabwe (1998)	Zimbabwe	297	35.8	ART codes and criteria for assessing restorations and sealants/criteria for diagnosing carious lesions in ART studies	95.7	92	85.3

Study title and year of publication	Country	Total procedures performedn	Sample loss %	Evaluation criteria	Survival rate after one year of follow up %	Survival rate after two years of follow up %	Survival rate after two to three years of follow up %
ART restorations and glass ionomer sealants in Zimbabwe: survival after 3 years (1998)	Zimbabwe	297	38.6	ART codes and criteria for assessing restorations and sealants/criteria for diagnosing carious lesions in ART studies	98.6	93.8	88.3
The implementation and preliminary evaluation of an ART strategy in Mexico: a country example (2009)	Mexico	314	10	ART codes and criteria for assessing restorations and sealants/criteria for diagnosing carious lesions in ART studies	80.2		
The atraumatic restorative treatment (ART) strategy in Mexico: two-years follow up of ART sealants and restorations (2013)	Mexico	291	0	ART codes and criteria for assessing restorations and sealants/criteria for diagnosing carious lesions in ART studies	82.1	74	
Atraumatic restorative treatment and glass ionomer sealants in Tunisian children: survival after 3 years (2002)	Tunisia	97	41.2	ART criteria	88.6	49.1	64.0
Studies that follow the ART restorations in single surfaces: Permanent Teeth							
The implementation and preliminary evaluation of an ART strategy in Mexico: a country example. (2009)	Mexico	96	8.4	ART codes and criteria for assessing restorations and sealants/criteria for diagnosing carious lesions in ART studies	87.5		
The atraumatic restorative treatment (ART) strategy in Mexico: two-years follow up of ART sealants and restorations. (2013)	Mexico	74	0	ART codes and criteria for assessing restorations and sealants / Criteria for diagnosing carious lesions in ART studies	86.5	80.9	
Atraumatic restorative treatment and glass ionomer sealants in Tunisian children: survival after 3 years (2002)	Tunisia	126	31.8	ART criteria	88.1	56.1	45.7

Source: research data.

There were also available data from ART sealants, but we chose not to include them, as it was not the purpose of this review.

The survival rates were very high, which indicates that ART, if correctly indicated and performed, can be routine in health services, regardless of electricity, HDI, or location. It is worth mentioning that the criteria used in the included studies evaluated the presence of complete restoration or wear that did not require repair, but in the context of large-scale implantation, the presence of restorative material with

failures that need repair can be considered a success, if there is no active caries lesion or recurrence. In addition, in cases of fracture or total loss of the restoration, if the carious lesion is paralyzed, it can also be a positive result because more invasive treatments are avoided. However, only follow-ups of occlusal cavities were performed by the researchers and in only two countries^{24,25,32-34}. The difficulty of monitoring patients in routine care is understood, especially in the context of implementation, in which professionals are still adapting to new practices. Still, efforts are needed to include

the monitoring of procedures performed on teeth with one or more tooth surfaces injured as routine in health services.

All countries increased access to health services after implementation^{16,25,29}, especially in municipalities with a low human development index (HDI), as in the case of Mexico^{24,25}, or in rural or remote areas, as in Bolivia¹⁹. However, although the treatment was designed to be conducted in territories where there was no access to conventional treatment, in countries where studies were conducted to evaluate the quality and longevity of restorations, such as Mexico²⁵ and Zimbabwe³²⁻³⁴, results were close to those of systematic reviews involving

controlled clinical trials.

3.3.3 Qualitative data

Regarding facilitators and barriers, four cross-sectional studies (questionnaires) assessed the perceptions of professionals and users of health services. Another seven studies (case studies, experience reports, and cohort) that contained data from the perceptions of patients and researchers were also included. Data identified by these studies as barriers or facilitators are presented in Table 2.

Table 2 - Qualitative data - Barriers and facilitators

Barriers		
	Studies N (%)	Countries where the Included Studies Report Barriers
Barriers perceived by health team professionals		
Finance	5 (83.3)	Cambodia, Mexico, Tanzania
Lack of supplies	4 (66.5)	Americas, Cambodia, Mexico, Tanzania
Lack of induction by managers	3 (50)	Americas, Tanzania
Technical ability	1 (16.6)	Tanzania
Resistance to change	2 (33.3)	Americas, Tanzania
Total	6 studies	
Barriers perceived by health service users		
Common sense that ART is of inferior quality	2 (100)	Americas, Tanzania
Lack of knowledge	1 (50)	Americas
Total	2 studies	
Barriers perceived by researchers		
Lack of supplies	3 (60)	South Africa, Cambodia, Mexico
Lack of human resources	2 (40)	South Africa, Cambodia
Common sense of health professionals	2 (40)	South Africa, Tunisia
Common sense of health service users	1 (20)	South Africa
Total	5 studies	
Facilitators		
	Studies N (%)	Countries where the included studies report facilitators
Facilitators perceived by health professionals		
Perception through the practice of ART in routine and positive experiences	3 (100)	Americas, Cambodia, Tanzania
Permanent education	2 (66.6)	Mexico, Cambodia
Coverage increase	1 (33.3)	Americas
Total	3 studies	
Facilitators perceived by health service users		
Favorable opinion after knowing about ART	1 (50)	Americas
Satisfaction with the treatment received	1 (50)	Bolivia
Treatment fear/anxiety reduction	1 (50)	Bolivia
Total	2 studies	
Facilitators perceived by researchers		
Coverage increase	1 (33.3)	Bolivia
Permanent Education	2 (66.6)	Bolivia, Cambodia
Have availability of necessary supplies in health services	1 (33.3)	Cambodia
Total	3 studies	

Source: research data.

Regarding the barriers identified in the implementation, the lack of supplies (dental instruments and restorative materials) to perform the procedures and induction by the service managers (through stipulation of targets, monitoring, financial resources) seems to have more influence than the other barriers mentioned.

One of the studies conducted in Cambodia²¹ pointed out that after years of offering well-structured courses to train dental nurses who would work in remote provinces in the country, little was done concerning restorative treatments. During the execution of the research, a health unit received the necessary inputs, received quarterly visits by members of the

Ministry of Health, and goals were agreed upon. At the same time, there was no interference in the other, which resulted in a significant increase in minimally invasive restorative treatments in this unit. Therefore, it seems that permanent education alone, although fundamental for updating and changing professionals' perceptions, is not enough for new practices to be established. Political instability, lack of support in institutions, and lack of engaged and participative leadership were identified as determining factors in cases where ART was not consolidated as a sustainable public policy²⁹.

In addition to the common barriers reported, it is worth highlighting an important finding in a study conducted in Bolivia, as information related to the dosage of components of glass ionomer cement for handling under extreme environmental conditions is not described in detail in technical profiles of manufacturers (in the case of Bolivia, altitudes well above sea level and relative air humidity ranging from 20% to more than 90%). This can be related to obtaining an adequate viscosity, resistance, and adherence of the products, which can negatively impact the quality of the results¹⁹. In the case of Bolivia, the material handling protocols were defined based on a consensus with a committee of experts.

As for the facilitators, the findings show that permanent education and professional practice enhance the use of ART as routine practice in health services and the availability of inputs and induction by managers, as already discussed. In addition, regarding service users, there seems to be high satisfaction with the approach and reduction of fear/anxiety in dental treatments¹⁹. The scientific literature corroborates these findings since the first clinical studies with ART reported greater comfort and acceptance of patients, especially children, as they do not receive dental anesthesia, rubber dam isolation, and use of bur. Another factor that contributes to acceptance is the possibility of performing treatments in schools, where children are, in general, less anxious than those treated in clinics or dental offices⁸.

Despite the high performance of the technique, that has been scientifically proven over the last decades, the implementation of ART as a policy has only been identified in developing countries, that are in most cases without universal health systems. Therefore, it was not possible to evaluate its implementation in better structured healthcare systems, which could guarantee conditions for the sustainability of ART with less structural difficulties.

Finally, at the beginning of 2020, the WHO declared the spread of COVID-19 to be pandemic. The COVID-19 has had a particular impact on dentistry. Studies indicate sites of the oral cavity as possible entries of the coronavirus and that angiotensin receptors present in salivary gland ducts can be the primary target of pathogenic cellular invasion^{44,45}. Moreover, as many dental procedures generate aerosols, such procedures and practices are being reconsidered. Faced with this scenario, aggravated by the scarcity of personal protective equipment, lack of clear biosafety protocols, the need to

preserve health teams and reduce risks of contamination by users, health systems around the world initially suspended elective procedures in dentistry, with urgent and emergency maintenance only, as recommended by several institutions and local governments⁴⁷⁻⁵¹.

In this context, Minimal Intervention Dentistry deserves even more attention from researchers, health service managers, and policymakers. It enables other types of health care that are not restricted to operative procedures³⁸. In addition, non-generating aerosol procedures, such as ART, are necessary for the resumption of oral health in the trans and post-pandemic periods. New policies and programs must be well-structured for developing and implementing ART in routine dental care.

4 Conclusions

There is sufficient scientific evidence for the use of ART as a modern and safe strategy for preventive and restorative care. In the context of public policies, the approach is still at an early stage and there seems to be a gap between knowledge of ART and its implementation. Therefore, efforts are needed to reduce this gap so that practices that meet the modern philosophy of health care, as is the case with ART, do not stay only on the political agenda globally, but also widespread, and integrated in public health systems.

References

1. Brasil. Ministério da Saúde. Síntese de Evidências para Políticas de Saúde: distribuição geográfica e fixação dos dentistas na atenção primária do Sistema Único de Saúde. Brasília: MS; 2022.
2. OMS - Organização Mundial da Saúde. Declaração de Alma-Ata. Conferência Internacional sobre cuidados de Saúde set 6-8 1978; Alma-Ata, URSS. Alma-Ata; 1978.
3. Tesch FC, De Oliveira BH, Leão A. Mensuração do impacto dos problemas bucais sobre a qualidade de vida de crianças: aspectos conceituais e metodológicos. *Cad Saúde Publ* 2007;23(11):2555-64.
4. GBD. Disease and Injury Incidence and Prevalence Collaborators Global, regional, and national incidence, prevalence, and years lived with disability for 328 diseases and injuries for 195 countries, 1990-2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet* 2017 2016;390:1211-159.
5. Frencken JE, Holmgren CJ, van Palenstein Helderma WH. Basic Package of Oral Care. [access in: 5 jun 2023]. Available in: <http://www.chdentalinstitute.org/images/BPOC.pdf>.
6. Frencken JO, Holmgren CJ. Atraumatic restorative treatment for dental caries. Nijmegen: STI. Book 1999
7. Frencken JE, Leal SC, Navarro MF. Twenty-five-year atraumatic restorative treatment (ART) approach: a comprehensive overview. *Clin Oral Investig* 2012;16(5):1337-46. doi: 10.1007/s00784-012-0783-4.
8. Navarro MFL, Leal SC, Molina GF, Sarmiento R. Tratamento restaurador atraumático: atualidades e perspectivas. *Rev Assoc Paul Cir Dent* 2015;69(3):289-301.
9. Mickenautsch S, Yengopal V, Banerjee A. Atraumatic restorative treatment versus amalgam restoration longevity: a

- systematic review. *Clin Oral Investig* 2010;14(3):233-40. doi: 10.1007/s00784-009-0335-8.
10. Tedesco TK, Bonifácio CC, Calvo AFB, Gimenez T, Braga MM, Raggio DP. Caries lesion prevention and arrestment in approximal surfaces in contact with glass ionomer cement restorations: a systematic review and meta-analysis. *Int J Paediatr Dent* 2016;26(3):161-72. doi: 10.1111/ipd.12174.
 11. Gauvin F-P. Understanding policy developments and choices through the “3-i” framework: interests, ideas and institutions. *Que: National Collaborating Centre for Health Public Policy*; 2014.
 12. Simpson DD. A Framework for implementing sustainable oral health promotion interventions. *J Public Health Dent* 2011;71(s1);Suppl 1:S84-94.
 13. Oliveira APC, Poz MRD, Craveiro I, Gabriel M, Dussault G. Fatores que influenciaram o processo de formulação de políticas de recursos humanos em saúde no Brasil e em Portugal: estudo de caso múltiplo. *Cad Saude Publica* 2018;34(2):e00220416. doi: <https://doi.org/10.1590/0102-311X00220416>.
 14. Souza C. Políticas públicas: conceitos, tipologias e sub-áreas. Salvador: Fundação Luís Eduardo Magalhães; 2012.
 15. Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ* 2021; 372:71. doi: <https://doi.org/10.1136/bmj.n71>
 16. Mickenautsch S, Rudolph MJ. Implementation of the ART approach in South Africa: an activity report. *SADJ* 2001;56(7):327-9.
 17. Mickenautsch S, Frencken JE. Utilization of the ART approach in a group of public oral health operators in South Africa: a 5-year longitudinal study. *BMC Oral Health* 2009;9:10. doi: 10.1186/1472-6831-9-10.
 18. Mickenautsch S, Frencken JE, Van't Hof M. Factors inhibiting the implementation of the Atraumatic Restorative Treatment approach in public oral health services in Gauteng Province, South Africa. *J Appl Oral Sci* 2007;15(1):1-8. doi: 10.1590/s1678-77572007000100002.
 19. Lopez J, Andrade G. El tratamiento restaurador atraumático (TRA) como parte de un paquete básico de servicios de salud oral en el área rural de Bolivia Evidencia sobre la efectividad de la aplicación del TRA y de acciones de promoción de la salud oral en escolares entre 6 y 14 años de edad, en áreas rurales con población dispersa. *Glob Health Promot* 2011;18(1):146-8. doi: <https://doi.org/10.1177/1757975910393201>.
 20. Mallow PK, Klaipo M, Durward CS. Dental nurse training in Cambodia - a new approach. *Int Dent J* 1997;47(3):148-56.
 21. Chher T, Hak S, Courtel F, Durward C. Improving the provision of the Basic Package of Oral Care (BPOC) in Cambodia. *Int Dent J* 2009;59(1):47-52.
 22. Farag A, van der Sanden WJM, Mulder J, Creugers NHJ, Frencken JE. Introduction of the ART approach in Egypt: intentions, clinical effects and perceived barriers. a cohort study. *Acta Odontol Scand* 2013;71(3/4):1014-121. doi: 10.3109/00016357.2012.741705.
 23. Farag AS. Atraumatic restorative treatment and oral health in Upper Egypt. Nijmegen: Radboud University Nijmegen; 2012.
 24. Hermosillo VH, Quintero LE, Guerrero ND. The implementation and preliminary evaluation of an ART strategy in Mexico: a country example. *J Appl Oral Sci* 2009;17;Suppl:114-21.
 25. Luengas-Quintero E, Frencken JE, Muñúzuri-Hernández JA. The atraumatic restorative treatment (ART) strategy in Mexico: two-years follow up of ART sealants and restorations. *BMC Oral Health* 2013;13:42. doi: 10.1186/1472-6831-13-42.
 26. Mandari GJ, Matee MIN. Atraumatic restorative treatment (ART): The Tanzanian experience. *Int Dent J* 2006;56(2):71-6.
 27. Kikwilu EN, Frencken J, Mulder J. Sciences A. Impact of Atraumatic Restorative Treatment (ART) on the treatment profile in pilot government dental clinics in Tanzania. *BMC Oral Health* 2009;9:14. doi: 10.1186/1472-6831-9-14.
 28. Kikwilu EN, Frencken JE, Mulder J. Barriers to the adoption of the ART approach as perceived by dental practitioners in governmental dental clinics, in Tanzania. *J Appl Oral Sci* 2009;17(5):408-13. doi: <https://doi.org/10.1590/S1678-77572009000500011>.
 29. Kikwilu EN, Frencken JE, Mulder J, Masalu JR, Joyce R. Dental practitioners' attitudes, subjective norms and intentions to practice atraumatic restorative treatment (ART) in Tanzania. *J Appl Oral Sci* 2009;17(2):97-102.
 30. Babo Soares LF, Bettiol SS, Dalla-Fontana IJ, et al. Opportunities in oral health policy for Timor-Leste. *WHO South East Asia J Public Health* 2016;5(2):164-73. doi: 10.4103/2224-3151.206254.
 31. Abid A, Chkir F, Salem K, Argoubi K, Sfar-Gandoura M. 2002. Atraumatic Restorative Treatment and glass ionomer sealants in Tunisian children: survival after 3 years. *East Mediterr Health J* 2002;8(2/3):315-23.
 32. Frencken JE, Makoni F, Sithole WD. Atraumatic restorative treatment and glass-ionomer sealants in a school oral health program in Zimbabwe: evaluation after 1 year. *Caries Res* 1996;30(6):428-33. doi: 10.1159/000262355.
 33. Frencken JE, Makoni F, Sithole WD, Hackenitz E. Three-year survival of one-surface ART restorations and glass-ionomer sealants in a school oral health programme in Zimbabwe. *Caries Res* 1998;32(2):119-26. doi: 10.1159/000016441.
 34. Frencken JE, Makoni F, Sithole WD. ART restorations and glass ionomer sealants in Zimbabwe: survival after 3 years. *Commun Dent Oral Epidemiol* 1998;26(6):372-81. doi: 10.1111/j.1600-0528.1998.tb01975.x.
 35. Ruiz O, Frencken JE. ART integration in oral health care systems in Latin American countries as perceived by directors of oral health. *J Appl Oral Sci* 2009;17;Suppl:106-13.
 36. Gonçalves CÂ, Vazquez FL, Ambrosano GMB, Mialhe FL, Pereira AC, Sarracini KLM, et al. Estratégias para o enfrentamento do absenteísmo em consultas odontológicas nas Unidades de Saúde da Família de um município de grande porte: uma pesquisa-ação. *Cienc Saúde Colet* 2015;20(2):449-60. doi: <http://dx.doi.org/10.1590/1413-81232015202.00702014>.
 37. Jandrey CM, Drehmer TM. Absenteísmo no atendimento clínicodontológico: o caso do módulo de saúde comunitária do CPOS. *Rev Fac Odontol Porto Alegre* 2000;40(2):24-8.
 38. Cruz DF, Cavalcanti RP, Lucena EHGd, Padilha WVN. A regulação da atenção à saúde bucal e o absenteísmo: cenários e possibilidades. *Rev Fam Ciclos Vida Saúde Contexto Soc* 2018;6(2):228-37. doi: 10.18554/refacs.v6i2.2820.
 39. Ericson D, Kidd E, McComb D, Mjör I, Noack MJ. Minimally invasive dentistry - concepts and techniques in cariology.

- Oral Health Prev Dent 2003;1(1):59-72.
40. Amara N, Ouimet M, Landry R. New evidence on instrumental, conceptual, and symbolic utilization of university research in government agencies. *Sci Commun* 2004;26(1):75-106.
 41. WHO Oral Health programme. Atraumatic restorative treatment (ART) for tooth decay: a global initiative. Geneva: World Health Organization; 1998.
 42. Barreto JOM, Souza NM. Avançando no uso de políticas e práticas de saúde informadas por evidências: a experiência de Piripiri-Piauí. *Cien Saúde Colet* 2013;18(1):25-34. doi: <http://dx.doi.org/10.1590/S1413-81232013000100004>.
 43. Lavis JN, Røttingen JA, Bosch-Capblanch X, et al. Guidance for evidence-informed policies about health systems: linking guidance development to policy development. *PLOS Med* 2012;9(3):e1001186. doi: <https://doi.org/10.1371/journal.pmed.1001186>.
 44. Dias RI, Barreto JOM, Souza NM. Desenvolvimento atual da Rede de Políticas Informadas por Evidências (EVIPNet Brasil): relato de caso. *Rev Panam Salud Publica* 2014;36(1):50-6.
 45. Liu L, Wei Q, Alvarez X. Epithelial cells lining salivary gland ducts are early target cells of severe acute respiratory syndrome coronavirus infection in the upper respiratory tracts of rhesus macaques. *J Virol* 2011;85(8):4025-30. doi: [10.1128/JVI.02292-10](https://doi.org/10.1128/JVI.02292-10).
 46. Matuck BF, Dolhnikoff M, Maia GVA, Sendyk DI, Zarpellon A, Costa Gomes S, et al. Periodontal tissues are targets for Sars-Cov-2: a post-mortem study. *J Oral Microbiol* 2021;13:1, 1848135, doi: [10.1080/20002297.2020.1848135](https://doi.org/10.1080/20002297.2020.1848135).
 47. Ministério da Saúde. Secretaria de Atenção Primária à Saúde (SAPS). Atendimento odontológico no SUS. Brasília: MS; 2020.
 48. Argentina: Ministério de La Salud Covid. 19 ATENCIÓN odontológica programada inicial. [access in 8 jun 2020]. Available in: <http://www.msal.gov.ar/images/stories/bes/graficos/0000001937cnt-covid-19recomendaciones-atencion-odontologica-programada.pdf>.5.
 49. Chile, Minsal Orientaciones para atención odontológica en Fase IV COVID-19. Available in: <https://diprece.minsal.cl/wp-content/uploads/2020/03/ORIENTACIONES-ATENCIONODONTOLOGICAS-COVID-19-.pdf>.6.
 50. American Dental Association, ADA. ADA recommending dentists postpone elective procedures. American Dental Association; 2020. [access in 8 jun 2022]. Available in: <https://www.ada.org/en/publications/ada-news/2020-archive/march/adarecommending-dentists-postpone-elective-procedures>.
 51. Kochhar AS, Bhasin R, Kochhar GK, Dadlani H. Provision of continuous dental care for oral oncology patients during & after COVID-19 pandemic. *Oral Oncol* 2020;106:104785. doi: [10.1016/j.oraloncology.2020.104785](https://doi.org/10.1016/j.oraloncology.2020.104785).